

JATENZO[®]
(testosterone undecanoate)
Capsules 

IT'S GO TIME.

HELPING YOU GET STARTED ON JATENZO[®]

JATENZO[®]
(testosterone undecanoate)
Capsules 

**PAY AS LITTLE AS \$0 FOR
YOUR MONTHLY PRESCRIPTION***

Use this for savings & treatment support

Provide information below to
pharmacy to receive copay offer

RxBIN: 610020

PCN: ACR

Group: 99995028

Identification: 42435701301



*Must have commercial insurance coverage for your prescription.

*See back cover for eligibility details and
other information about savings information

GetJATENZO.com

What is JATENZO?

JATENZO is approved by the FDA as a prescription drug used to treat adult men who have low or no testosterone levels due to certain medical conditions.

JATENZO is a controlled substance (CIII) because it contains testosterone. It is not known if JATENZO is safe or effective in children younger than 18 years old. Improper use of JATENZO may affect bone growth in children.

IMPORTANT SAFETY INFORMATION FOR JATENZO (testosterone undecanoate)

What is the most important information I should know about JATENZO?

JATENZO can cause serious side effects, including:

- JATENZO can increase your blood pressure, which can increase your risk of having a heart attack or stroke and can increase your risk of death due to a heart attack or stroke. Your risk may be greater if you have already had a heart attack or stroke or if you have other risk factors for heart attack or stroke.
- If your blood pressure increases while on JATENZO, blood pressure medicines may need to be started. If you are taking blood pressure medicines, new blood pressure medicines may need to be added or your current blood pressure medicines may need to be changed to control your blood pressure.
- If your blood pressure cannot be controlled, JATENZO may need to be stopped.
- Your healthcare provider will monitor your blood pressure while you are being treated with JATENZO.

Please see full Important Safety Information on pages 10-11, and visit Jatenzo.com for full Prescribing Information, including BOXED WARNING on increases in blood pressure, and Medication Guide for JATENZO.



JATENZO
(testosterone undecanoate)
Capsules 

UNDERSTAND YOUR DIAGNOSIS

When your doctor tests your testosterone levels and finds low or absence of testosterone due to certain medical conditions, this is called hypogonadism.

You're not alone, though. This condition affects 30% of men in the United States*. Seeking out treatment is understandable. **By working with your doctor, you're already taking the right steps.**

There are several options to treat your condition, including (but not limited to):

- JATENZO® (an oral softgel)
- Patches
- Gel
- Implanted pellets
- Injection

It's important to talk to your doctor to find out which option may be right for you.

SELECT IMPORTANT SAFETY INFORMATION

What is JATENZO?

JATENZO is a prescription medicine that contains testosterone. JATENZO is used to treat adult men who have low or no testosterone due to certain medical conditions.

JATENZO is a controlled substance (CIII) because it contains testosterone that can be a target for people who abuse prescription medicines. Keep your JATENZO in a safe place to protect it. Never give your JATENZO to anyone else, even if they have the same symptoms you have. Selling or giving away this medicine may harm others and is against the law.

JATENZO is not meant for use by women.

Please visit Jatenzo.com for full Prescribing Information, including BOXED WARNING on increases in blood pressure, and Medication Guide for JATENZO.

*Peterson MD et al. *Nature.com/Scientific Reports*. (2018) 8:5897; 5-9



TESTOSTERONE
REPLACEMENT
IN AN ORAL SOFTGEL
IS ALL RIGHT HERE.

IT'S FDA-APPROVED AND
IT'S NOT STUCK ON,
SMEARED AROUND,
OR INJECTED.

WELCOME TO THE JATENZO[™] ZONE

DIAL IT IN

JATENZO[®]
(testosterone undecanoate)
Capsules 

JATENZO[®] is the first oral softgel that offers you **dosing flexibility** to fit your specific needs. All doses must be taken twice daily with food. Your doctor will check your testosterone levels from time to time to see how you respond to treatment and to see if any adjustments are needed.

Schedule a follow-up visit with your doctor a month after treatment begins. During your appointment, you can check your levels to make sure you're staying right where your doctor wants you to be.



JATENZO is the first oral softgel for low testosterone that offers flexible dosing to fit your needs.

YOUR SAFETY IS OUR PRIORITY



In clinical trials, **no liver toxicity-related side effects*** were observed with JATENZO. That's a fact that's important to us and may be good for you and your doctor to keep in mind when exploring treatment options.

The most common side effects reported with JATENZO in clinical studies included headache, increased volume percentage of red blood cells in your blood, and high blood pressure. See important safety information.

SELECT IMPORTANT SAFETY INFORMATION

Do not take JATENZO if you:

- have breast cancer.
- have or might have prostate cancer.
- are a woman who is pregnant. JATENZO may harm your unborn baby.
- are allergic to JATENZO or any ingredients in JATENZO.
- have low testosterone without certain medical conditions. For example, do not take JATENZO if you have low testosterone due to age.

Before you take JATENZO, tell your healthcare provider about all of your medical conditions, including if you:

- have high blood pressure or are treated for high blood pressure.
- have heart problems.
- have high red blood cell count (hematocrit) or high hemoglobin laboratory value.
- have urinary problems due to an enlarged prostate.
- have liver or kidney problems.
- have a history of mental health illness including suicidal thoughts or actions, depression, anxiety or mood disorder.
- have problems breathing while you sleep (sleep apnea).

Please visit Jatenzo.com for full **Prescribing Information**, including **BOXED WARNING on increases in blood pressure, and Medication Guide for JATENZO**.

*Statement true at the time of printing. Information is subject to change.



TAKE THE NEXT STEP

After you've been prescribed JATENZO, it's important to talk openly and honestly with your doctor about your treatment goals. From there, you can both work together on a plan to help you reach those goals.



TAKING YOUR DOSES

To achieve and maintain steady testosterone levels, each JATENZO® dose should be taken twice daily as prescribed by your doctor.



Once in the AM



Once in the PM

Remember to take each JATENZO dose with food as prescribed by your doctor.

JATENZO can be taken at home or on the go.

SELECT IMPORTANT SAFETY INFORMATION

Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements.

Using JATENZO with certain other medicines can affect each other. Especially, tell your healthcare provider if you take:

- insulin
- medicines that decrease blood clotting (blood thinners)
- corticosteroids
- medicines that increase blood pressure such as some cold medicines and pain medicines

Know the medicines you take. Ask your healthcare provider or pharmacist for a list of these medicines, if you are not sure. Keep a list of them and show it to your healthcare provider and pharmacist when you get a new medicine.

Please visit Jatenzo.com for full Prescribing Information, including BOXED WARNING on increases in blood pressure, and Medication Guide for JATENZO.

CONVERSATIONS ARE THE KEY

Set up a 30-day follow-up with your doctor after treatment begins. That way, you can take a look at any adjustments that may need to be made.

JATENZO is an FDA-approved testosterone replacement therapy with a safe oral delivery to help raise testosterone levels.

To learn more about JATENZO, visit GetJATENZO.com



NOTES WORTH TAKING

As you start treatment, take notes on how you're feeling and what dose you're on. If your doctor adjusts your dose, look back on these notes to discuss how you're responding to treatment.

HOW ARE YOU FEELING TODAY?

What 3 symptoms of hypogonadism concern you most
(On a scale of 1-5: 5=very concerned; 1=not concerned at all)

1. 1 2 3 4 5
2. 1 2 3 4 5
3. 1 2 3 4 5

WHAT WAS YOUR TESTOSTERONE LEVEL FROM YOUR LAST DOCTOR'S VISIT?

It's good to know your baseline – you can use it to monitor your treatment progress.

..... ng/dL.

HOW OFTEN ARE YOU TAKING YOUR TREATMENT?

WHAT'S YOUR STARTING DOSE OF JATENZO?

..... mg, twice daily with food

WHEN IS YOUR FOLLOW-UP APPOINTMENT?

.....

FOLLOW-UP NOTES

After your follow-up appointment, come back to these pages to talk with your doctor about how you're responding to treatment. If you've used the Doctor Discussion Guide on GetJATENZO.com, you can also share that information with your doctor during this time.

HOW ARE YOU FEELING TODAY?

How do you feel about your symptoms today?
(On scale of 1-5: 5=much better; 1=much worse)

1. 1 2 3 4 5
2. 1 2 3 4 5
3. 1 2 3 4 5

DO YOU HAVE ANY CONCERNS?

WHAT DIFFERENCES – IF ANY – HAVE YOU NOTICED SINCE YOUR LAST VISIT WITH YOUR DOCTOR?

.....
.....

WHEN LAST MEASURED, WERE YOUR TESTOSTERONE LEVELS LOW OR IN THE NORMAL RANGE?

Normal levels are typically in the 300-1000 ng/dL range.

WHAT'S YOUR NEW DAILY DOSE?

..... mg, twice daily with food.

SCAN THIS CODE WITH
YOUR PHONE CAMERA



to visit the FAQs section on [GetJATENZO.com](https://www.getjatenzo.com).

There, you'll find answers to commonly asked questions about JATENZO®.

What are JATENZO® capsules approved for?

JATENZO is approved by the FDA as a prescription drug used to treat adult men who have low or no testosterone levels due to certain medical conditions.

JATENZO is a controlled substance (CIII) because it contains testosterone. It is not known if JATENZO is safe or effective in children younger than 18 years old. Improper use of JATENZO may affect bone growth in children.

IMPORTANT SAFETY INFORMATION FOR JATENZO (testosterone undecanoate)

What is the most important information I should know about JATENZO?

JATENZO can cause serious side effects, including:

- **JATENZO can increase your blood pressure, which can increase your risk of having a heart attack or stroke and can increase your risk of death due to a heart attack or stroke. Your risk may be greater if you have already had a heart attack or stroke or if you have other risk factors for heart attack or stroke.**
- **If your blood pressure increases while on JATENZO, blood pressure medicines may need to be started. If you are taking blood pressure medicines, new blood pressure medicines may need to be added or your current blood pressure medicines may need to be changed to control your blood pressure.**
- **If your blood pressure cannot be controlled, JATENZO may need to be stopped.**
- **Your healthcare provider will monitor your blood pressure while you are being treated with JATENZO.**

What is JATENZO?

JATENZO is a prescription medicine that contains testosterone. JATENZO is used to treat adult men who have low or no testosterone due to certain medical conditions.

JATENZO is a controlled substance (CIII) because it contains testosterone that can be a target for people who abuse prescription medicines. Keep your JATENZO in a safe place to protect it. Never give your JATENZO to anyone else, even if they have the same symptoms you have. Selling or giving away this medicine may harm others and is against the law.

JATENZO is not meant for use by women.

Do not take JATENZO if you:

- have breast cancer.
- have or might have prostate cancer.
- are a woman who is pregnant. JATENZO may harm your unborn baby.
- are allergic to JATENZO or any ingredients in JATENZO.
- have low testosterone without certain medical conditions. For example, do not take JATENZO if you have low testosterone due to age.

Before you take JATENZO, tell your healthcare provider about all of your medical conditions, including if you:

- have high blood pressure or are treated for high blood pressure.
- have heart problems.
- have high red blood cell count (hematocrit) or high hemoglobin laboratory value.
- have urinary problems due to an enlarged prostate.
- have liver or kidney problems.
- have a history of mental health illness including suicidal thoughts or actions, depression, anxiety or mood disorder.
- have problems breathing while you sleep (sleep apnea).

Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements.

Using JATENZO with certain other medicines can affect each other. Especially, tell your healthcare provider if you take:

- insulin
- medicines that decrease blood clotting (blood thinners)
- corticosteroids
- medicines that increase blood pressure such as some cold medicines and pain medicines

Know the medicines you take. Ask your healthcare provider or pharmacist for a list of these medicines, if you are not sure. Keep a list of them and show it to your healthcare provider and pharmacist when you get a new medicine.

How should I take JATENZO?

- Take JATENZO exactly as your healthcare provider tells you to take it.
- Take JATENZO by mouth two times daily. Take 1 time in the morning and take 1 time in the evening.
- Take JATENZO with food.
- Your healthcare provider may change your JATENZO dose. Do not change your JATENZO dose without talking to your healthcare provider.

What are the possible side effects of JATENZO?

JATENZO may cause serious side effects including:

- See “**What is the most important information I should know about JATENZO?**”
- **Increase in blood pressure.**
 - JATENZO can increase your blood pressure, which can increase your risk of having a heart attack or stroke and can increase your risk of death due to heart attack or stroke. Your risk may be greater if you already have heart disease or if you have had a heart attack or stroke.
 - You may need to start new medicines or have medicines changed for high blood pressure while on JATENZO.
 - Your healthcare provider should check your blood pressure while you use JATENZO.
- **Increase in red blood cell count (hematocrit) or hemoglobin.**
 - JATENZO increases red blood cell counts in some patients. High red blood cell counts increase the risk of blood clots, strokes, and heart attacks.
 - You may need to stop JATENZO if your red blood cell count increases.
 - Your healthcare provider should check your red blood cell count and hemoglobin while you use JATENZO.
- **If you already have an enlarged prostate, your signs and symptoms may worsen while using JATENZO.** These may include:
 - increased urination at night
 - trouble starting your urine stream
 - urinating many times during the day
 - urge to go to the bathroom right away
 - a urine accident
 - inability to pass urine or weak urine flow
- **Increased risk of prostate cancer.** Your healthcare provider should check you for prostate cancer or any other prostate problems before you start and while you use JATENZO.
- **Blood clots in the legs or lungs.** Signs and symptoms of a blood clot in your leg can include pain, swelling or redness. Signs and symptoms of a blood clot in your lungs can include difficulty breathing or chest pain.
- **Abuse.** Testosterone can be abused, when taken at higher than prescribed doses and when used with other anabolic androgenic steroids. Abuse can cause serious heart and psychological side effects. Your healthcare provider should check you for signs of abuse before and during treatment with JATENZO.
- **In large doses JATENZO may lower your sperm count.**

• **Liver problems.** Symptoms of liver problems may include:

- nausea or vomiting
- yellowing of your skin or whites of your eyes
- dark urine
- pain on the right side of your stomach area (abdominal pain)

• **Swelling of your ankles, feet, or body (edema), with or without heart failure.**

• **Enlarged or painful breasts.**

• **Breathing problems while you sleep (sleep apnea).**

• **Changes in mood.** Talk to your healthcare provider if you have changes in mood or behavior including, new or worsening depression, or suicidal thoughts.

Call your healthcare provider right away if you have any of the serious side effects listed above.

The most common side effects of JATENZO include:

- increased red blood cell count
- burping
- enlarged prostate gland
- diarrhea
- foot, leg, and ankle swelling
- high blood pressure
- heart burn
- nausea
- headache

Other side effects include more erections than are normal for you or erections that last a long time.

Tell your healthcare provider if you have any side effect that bothers you or that does not go away.


These are not all the possible side effects of JATENZO. For more information, ask your healthcare provider or pharmacist. Call your doctor for medical advice about side effects.

Keep JATENZO and all medicines out of the reach of children.

Please visit Jatenzo.com for full Prescribing Information, including BOXED WARNING on increases in blood pressure, and Medication Guide for JATENZO.

You are encouraged to report any side effects of prescription drugs to the FDA. Visit www.fda.gov/medwatch or call 1-800-FDA-1088.

JATENZO®
(testosterone undecanoate)
Capsules @



Save big on your monthly JATENZO® prescription*

If you're an eligible patient with commercial insurance, you may pay as little as \$0 for your monthly prescription using one of our network pharmacies. If your commercial insurance rejects JATENZO coverage, you will only pay \$150 for your prescription. If you do not have commercial insurance, you may purchase your JATENZO for \$185.

*See eligibility restrictions below.

Ask your doctor to send your JATENZO prescription through one of our network pharmacies to receive additional co-pay assistance.

TERMS, CONDITIONS & RESTRICTIONS

Eligible patients with commercial insurance will pay as little as \$0 for their monthly prescriptions, up to a maximum benefit amount applied per calendar year. Eligible patients without commercial insurance may still save up to 50% on their monthly prescription, up to a maximum benefit amount applied per calendar year. Offer valid for up to 12 uses. A valid Prescriber ID# is required on the prescription.

Patient Instructions: To redeem this offer, you must have a valid prescription for JATENZO. Follow the dosage instructions given by the doctor. This offer may not be redeemed for cash. By using this offer, you are certifying that you meet the eligibility criteria and will comply with the terms and conditions described in the Restrictions section below. Patients with questions about accessing the JATENZO Savings Card offer should ask their pharmacist to call the PDMI online processing Help Desk at 316-219-4802.

Pharmacist: When you apply this offer, you are certifying that you have not submitted a claim for reimbursement under any federal, state, or other governmental programs for this prescription. Participation in this program must comply with all applicable laws and regulations as a pharmacy provider. By participating in this program, you are certifying that you will comply with the terms and conditions described in the Restrictions section below.

Pharmacist Instructions for a Patient with an Eligible Third Party: Submit the claim to the primary Third-Party Payer first, then submit the balance due to PDMI as a Secondary Payer COB (coordination of benefits) with patient responsibility amount and a valid Other Coverage Code, (e.g., 8). The patient is responsible for as little as \$0 for their monthly prescriptions, up to a maximum benefit amount applied per calendar year. Offer valid for up to 12 uses. Reimbursement will be received from PDMI. Valid Other Coverage Code required. For any questions regarding PDMI online processing, please call the Help Desk at **316-219-4802**.

Restrictions: This offer is valid in the United States and Puerto Rico. Offer not valid for prescriptions reimbursed under Medicaid, a Medicare drug benefit plan, TRICARE, or other federal or state health programs (such as medical assistance programs). Cash Discount Cards and other noninsurance plans are not valid as primary under this offer. If the patient is eligible for drug benefits under any such program, the patient cannot use this offer. By using this offer, the patient certifies that he will comply with any terms of his health insurance contract requiring notification to his payer of the existence and/or value of this offer. Offer not valid for patients under 18 years of age. It is illegal to (or offer to) sell, purchase, or trade this offer. This offer is not transferable and is limited to one offer per person. Not valid if reproduced. Void where prohibited by law. Tolmar reserves the right to rescind, revoke, or amend this offer without notice at any time.

Please see full Important Safety Information on pages 10-11, and visit Jatenzo.com for full Prescribing Information, including BOXED WARNING on increases in blood pressure, and Medication Guide for JATENZO.



© 2023 Tolmar, Inc. All rights reserved. Tolmar, JATENZO, The JATENZone and their associated logos are trademarks of th Tolmar Group. TPI.2022.3441.v3 03/23